

Complementarity of the Maldacena and Randall-Sundrum Pictures*

M. J. Duff[†] and James T. Liu[‡]

*Randall Laboratory, Department of Physics, University of Michigan,
Ann Arbor, MI 48109-1120*

Abstract

We revive an old result, that one-loop corrections to the graviton propagator induce $1/r^3$ corrections to the Newtonian gravitational potential, and compute the coefficient due to closed loops of the $U(N)$ $\mathcal{N} = 4$ super-Yang-Mills theory that arises in Maldacena's AdS/CFT correspondence. We find exact agreement with the coefficient appearing in the Randall-Sundrum brane-world proposal. This provides more evidence for the complementarity of the two pictures.

Typeset using REVTeX

*Research supported in part by NSF Grant PHY-9722090.

[†]mduff@umich.edu

[‡]jimliu@umich.edu

It is an old, and seemingly forgotten, result that one-loop corrections to the graviton propagator induce $1/r^3$ corrections to the gravitational potential [1,2]:

$$V(r) = \frac{Gm_1m_2}{r} \left(1 + \frac{\alpha G}{r^2} \right), \quad (1)$$

where G is the four-dimensional Newton's constant, $\hbar = c = 1$ and α is a purely numerical coefficient given, in the case of spins $s \leq 1$, by $45\pi\alpha = 12N_1 + 3N_{1/2} + N_0$, where N_s are the numbers of particle species of spin s going around the loop [3,4,5]. However, the importance of this result has recently become apparent in attempts [6,7,8,9,10] to relate two topical but, at first sight, different developments in quantum gravity. These are Maldacena's AdS/CFT correspondence [11,12,13] and the Randall-Sundrum brane-world mechanism [14].

The AdS/CFT correspondence in general relates the gravitational dynamics of a $(d+1)$ -dimensional anti-de Sitter spacetime, AdS_{d+1} , to a d -dimensional conformal field theory, CFT_d . In the case of $d = 4$, Maldacena's conjecture, based on the decoupling limit of D3-branes in Type *IIB* string theory compactified on S^5 , then relates the dynamics of AdS_5 to an $\mathcal{N} = 4$ superconformal $U(N)$ Yang-Mills theory on its four-dimensional boundary [11]. Other compactifications are also possible, leading to different SCFT's on the boundary. We note that, by choosing Poincaré coordinates on AdS_5 , the metric may be written as

$$ds^2 = e^{-2y/L} (dx^\mu)^2 + dy^2, \quad (2)$$

in which case the superconformal Yang-Mills theory is taken to reside at the boundary $y \rightarrow -\infty$.

The Randall-Sundrum mechanism, on the other hand, was originally motivated, not via the decoupling of gravity from D3-branes, but rather as a possible mechanism for evading Kaluza-Klein compactification by localizing gravity in the presence of an uncompactified extra dimension. This was accomplished by inserting a positive tension 3-brane (representing our spacetime) into AdS_5 . In terms of the Poincaré patch of AdS_5 given above, this corresponds to removing the region $y < 0$, and either joining on a second partial copy of AdS_5 , or leaving the brane at the end of a single patch of AdS_5 . In either case the resulting Randall-Sundrum metric is given by

$$ds^2 = e^{-2|y|/L} (dx^\mu)^2 + dy^2, \quad (3)$$

where $y \in (-\infty, \infty)$ or $y \in [0, \infty)$ for a 'two-sided' or 'one-sided' Randall-Sundrum brane respectively.

The similarity of these two scenarios led to the notion that they are in fact closely tied together. To make this connection clear, consider the one-sided Randall-Sundrum brane. By introducing a boundary in AdS_5 at $y = 0$, this model is conjectured to be dual to a cutoff CFT coupled to gravity, with $y = 0$, the location of the Randall-Sundrum brane, providing the UV cutoff. This extended version of the Maldacena conjecture [15] then reduces to the standard AdS/CFT duality as the boundary is pushed off to $y \rightarrow -\infty$, whereupon the cutoff is removed and gravity becomes completely decoupled. Note in particular that this connection involves a single CFT at the boundary of a single patch of AdS_5 . For the case of a brane sitting between two patches of AdS_5 , one would instead require two copies of the CFT, one for each of the patches.

It has been suggested [7,8,9] that a crucial test of this Randall-Sundrum version of the Maldacena conjecture would be to compare the $1/r^3$ corrections to Newton's law in both pictures. From the above, we see that the contribution of a single CFT, with $(N_1, N_{1/2}, N_0) = (N^2, 4N^2, 6N^2)$, is

$$V(r) = \frac{Gm_1m_2}{r} \left(1 + \frac{2N^2G}{3\pi r^2} \right). \quad (4)$$

Using the AdS/CFT relation $N^2 = \pi L^3/2G_5$ [11] and the one-sided brane-world relation $G = 2G_5/L$ [14,8], where G_5 is the five-dimensional Newton's constant and L is the radius of AdS_5 , this becomes

$$V(r) = \frac{Gm_1m_2}{r} \left(1 + \frac{2L^2}{3r^2} \right). \quad (5)$$

The coefficient of the $1/r^3$ term is $2/3$ of the Randall-Sundrum result quoted in [14], but in fact agrees with the more thorough analysis of [16]. We shall confirm below that a more careful analysis of the Randall-Sundrum picture using the results of [9,17] yields exactly the same answer as the above AdS/CFT calculation, thus providing strong evidence for the conjectured duality of the two pictures.

First we derive (4) in more detail by computing the lowest order quantum corrections to solutions of Einstein's equations. Working with linearized gravity, we begin by writing the metric as

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}, \quad (6)$$

so that

$$\sqrt{-g}g^{\mu\nu} \equiv \tilde{g}^{\mu\nu} = \eta^{\mu\nu} - \tilde{h}^{\mu\nu} + \dots, \quad (7)$$

where

$$\tilde{h}_{\mu\nu} = h_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}h^\alpha{}_\alpha. \quad (8)$$

In harmonic gauge, $\partial_\mu \tilde{g}^{\mu\nu} = 0$ (*i.e.* $\partial_\mu \tilde{h}^{\mu\nu} = 0$), the classical linearized Einstein equation reads

$$\square \tilde{h}_{\mu\nu}^c(x) = -16\pi G T_{\mu\nu}(x), \quad (9)$$

where the superscript c denotes the classical contribution. Fourier transforming to momentum space results in

$$\tilde{h}_{\mu\nu}^c(p) = -16\pi G \Delta_4(p) T_{\mu\nu}(p), \quad (10)$$

where $\Delta_4(p) = -1/p^2$ is the four-dimensional massless scalar propagator.

Incorporating one-loop corrections, the quantum corrected metric becomes

$$\tilde{h}_{\mu\nu} = \tilde{h}_{\mu\nu}^c + \tilde{h}_{\mu\nu}^q, \quad (11)$$

where the quantum correction $\tilde{h}_q^{\mu\nu}$ is given in momentum space by

$$\tilde{h}_q^{\mu\nu}(p) = D^{\mu\nu\alpha\beta}(p)\Pi_{\alpha\beta\gamma\delta}(p)\tilde{h}_c^{\gamma\delta}(p). \quad (12)$$

$D^{\mu\nu\alpha\beta}$ is the graviton propagator,

$$D^{\mu\nu\alpha\beta}(p) = \frac{1}{2}\Delta_4(p)(\eta^{\mu\alpha}\eta^{\nu\beta} + \eta^{\mu\beta}\eta^{\nu\alpha} - \eta^{\mu\nu}\eta^{\alpha\beta} + \dots), \quad (13)$$

and $\Pi_{\alpha\beta\gamma\delta}$ is the one-loop graviton self-energy, which by symmetry and Lorentz invariance must be of the general form

$$\begin{aligned} \Pi_{\alpha\beta\gamma\delta}(p) = p^4 & \left[\Pi_1(p^2)\eta_{\alpha\beta}\eta_{\gamma\delta} + \Pi_2(p^2)(\eta_{\alpha\gamma}\eta_{\beta\delta} + \eta_{\alpha\delta}\eta_{\beta\gamma}) + \Pi_3(p^2)(\eta_{\alpha\beta}\hat{p}_\gamma\hat{p}_\delta + \eta_{\gamma\delta}\hat{p}_\alpha\hat{p}_\beta) \right. \\ & \left. + \Pi_4(p^2)(\eta_{\alpha\gamma}\hat{p}_\beta\hat{p}_\delta + \eta_{\alpha\delta}\hat{p}_\beta\hat{p}_\gamma + \eta_{\beta\gamma}\hat{p}_\alpha\hat{p}_\delta + \eta_{\beta\delta}\hat{p}_\alpha\hat{p}_\gamma) + \Pi_5(p^2)\hat{p}_\alpha\hat{p}_\beta\hat{p}_\gamma\hat{p}_\delta \right]. \end{aligned} \quad (14)$$

The ellipses in (13) refer to gauge dependent terms in the propagator which make no contribution if coupled to conserved sources. Combining (12), (13) and (14), one thus obtains the quantum corrected metric in the form

$$h_{\mu\nu}^q(p) = -p^2 \left[2\Pi_2(p)\delta_\mu^\alpha\delta_\nu^\beta + \Pi_1(p)\eta_{\mu\nu}\eta^{\alpha\beta} + (\Pi_3(p) + \dots)\hat{p}_\mu\hat{p}_\nu\eta^{\alpha\beta} \right] \tilde{h}_{\alpha\beta}^c, \quad (15)$$

where non-physical gauge-dependent terms have again been dropped. Finally, combining both classical and one-loop quantum results at the linearized level yields

$$\begin{aligned} h_{\mu\nu}(p) = & -16\pi G\Delta_4(p)[T_{\mu\nu}(p) - \tfrac{1}{2}\eta_{\mu\nu}T^\alpha{}_\alpha(p)] \\ & -16\pi G[2\Pi_2(p)T_{\mu\nu}(p) + \Pi_1(p)\eta_{\mu\nu}T^\alpha{}_\alpha(p)]. \end{aligned} \quad (16)$$

Note that we have ignored the gauge-dependent term in $h_{\mu\nu}$ proportional to $\hat{p}_\mu\hat{p}_\nu$. It makes no contribution when $h_{\mu\nu}$ is attached to a conserved source $T_{\mu\nu}$ satisfying $p^\mu T_{\mu\nu} = p^\nu T_{\mu\nu} = 0$.

The actual form of the one-loop Π_i 's depend on the theory at hand. However for any massless theory in four-dimensions, after cancelling the infinities with the appropriate counterterms, the finite remainder must necessarily have the form

$$\Pi_i(p) = 32\pi G \left(a_i \ln \frac{p^2}{\mu^2} + b_i \right), \quad (17)$$

where a_i and b_i , ($i = 1, 2, 3, 4, 5$), are numerical coefficients and μ is an arbitrary subtraction constant having the dimensions of mass. In order to make connection with the Newtonian potential, we Fourier transform (16) back to coordinate space. For the static potential we obtain the expected $1/r$ behavior at the classical level, while the quantum term generates the claimed $1/r^3$ correction. In addition, the constant parts in (17) give rise to a regulator-dependent $\delta^3(\mathbf{r})$ contact interaction. However we have no real expectation that this one-loop perturbative result remains valid when continued down to zero size. Moreover, possible $r^{-3} \ln \mu r$ terms come only from the $\hat{p}_\mu\hat{p}_\nu$ terms in (15) and hence drop out. For a point source, $T_{00}(x) = m\delta^3(\mathbf{r})$, we obtain to this order

$$\begin{aligned} g_{00} = & -\left(1 - \frac{2Gm}{r} - \frac{2\alpha G^2 m}{r^3} \right), \\ g_{ij} = & \left(1 + \frac{2Gm}{r} + \frac{2\beta G^2 m}{r^3} \right) \delta_{ij}, \end{aligned} \quad (18)$$

where, in agreement with [2], $\alpha = 4 \cdot 32\pi(a_1 + 2a_2)$ and $\beta = -4 \cdot 32\pi a_1$. This yields the potential given in (1). Explicit calculations of the self-energy (17) for spin 1 [3], spin 1/2 (two-component fermions) [4] and (real conformally coupled) spin 0 [5] yield¹

$$\begin{aligned} a_i(s=1) &= 4a_i(s=1/2) = 12a_i(s=0) \\ &= \frac{1}{120(4\pi)^2}(-2, 3, 2, -3, 4). \end{aligned} \quad (19)$$

Note that all spins contribute with the same sign as they must by general positivity arguments on the self-energy [4]. Thus

$$\alpha = 2\beta = \frac{1}{45\pi}(12N_1 + 3N_{1/2} + N_0) = \frac{2N^2}{3\pi}, \quad (20)$$

as quoted in the introductory paragraph above.

This α coefficient also determines that part of the Weyl anomaly [18,19] involving the square of the Weyl tensor [20,21]:

$$g_{\mu\nu}\langle T^{\mu\nu} \rangle = b \left(F + \frac{2}{3} \square R \right) + b' G, \quad (21)$$

where

$$\begin{aligned} F &= C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma} = R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 2R_{\mu\nu} R^{\mu\nu} + R^2, \\ G &= *R_{\mu\nu\rho\sigma} *R^{\mu\nu\rho\sigma} = R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 4R_{\mu\nu} R^{\mu\nu} + \frac{1}{3} R^2, \end{aligned} \quad (22)$$

and where b and b' are constants

$$\begin{aligned} b &= \frac{1}{120(4\pi)^2} [12N_1 + 3N_{1/2} + N_0], \\ b' &= -\frac{1}{720(4\pi)^2} [124N_1 + 11N_{1/2} + 2N_0]. \end{aligned} \quad (23)$$

Note that for the $\mathcal{N} = 4$ SCFT, the coefficient of the (Riemann)² term, $b + b'$, vanishes [21]. The same result is obtained if one calculates the holographic Weyl anomaly using the AdS/CFT correspondence [22]. Thus $b = 3\alpha/128\pi = c/(4\pi)^2$, where the c is the central charge given in the normalization of [8]. For the central charge, one obtains $c = \pi L^3/8G_5$ [22], so that

$$G\alpha = \frac{GL^3}{3G_5} = \frac{2L^2}{3}, \quad (24)$$

where the second equality makes use of the brane-world relation $G = 2G_5/L$. Although we have focused on the $\mathcal{N} = 4$ SCFT to relate the coefficient appearing in Newton's law to

¹Note that a symmetry factor of 1/2 was omitted in Ref. [3]; this was subsequently corrected in Ref. [5]

the central charge, the result (24) is universal, being independent of which particular CFT appears in the AdS/CFT correspondence, which is just as well since the Randall-Sundrum coefficient does not depend on the details of the fields propagating on the brane.

We now turn to this brane-world, where the five-dimensional action has the form [14]

$$S = \int d^5x \sqrt{-g_{(5)}} [M^3 R_{(5)} - \Lambda] + \int d^4x \sqrt{-g_{(4)}} \mathcal{L}_{\text{brane}}. \quad (25)$$

Here M is the five-dimensional Planck mass, $M^3 = 1/(16\pi G_5)$, and Λ is the cosmological constant in the bulk. Small fluctuations of the metric on the brane may be represented by [14,9]

$$ds^2 = e^{-2|y|/L} [\eta_{\mu\nu} + h_{\mu\nu}(x, y)] dx^\mu dx^\nu + dy^2, \quad (26)$$

where L is the ‘radius’ of AdS,

$$R_{MNPQ}^{(5)} = -\frac{1}{L^2} (g_{MP}^{(5)} g_{NQ}^{(5)} - g_{MQ}^{(5)} g_{NP}^{(5)}), \quad (27)$$

and is related to Λ by $\Lambda = -12M^3/L^2$. The brane-world geometry has been chosen such that x^μ are coordinates along the 3-brane, while y is the coordinate perpendicular to the brane (which sits at $y = 0$).

Both brane and bulk quantities are contained in the linearized metric $h_{\mu\nu}(x, y)$. However, for comparison with the CFT on the brane, we are only concerned with the former. Hence we consider a matter source on the brane, and examine $h_{\mu\nu}(x) \equiv h_{\mu\nu}(x, y = 0)$. For this case, the results of [9,17] indicate

$$\begin{aligned} h_{\mu\nu}(p) = & -\frac{2}{LM^3} \Delta_4(p) [T_{\mu\nu}(p) - \frac{1}{2} \eta_{\mu\nu} T^\alpha{}_\alpha(p)] \\ & -\frac{1}{M^3} \Delta_{KK}(p) [T_{\mu\nu}(p) - \frac{1}{3} \eta_{\mu\nu} T^\alpha{}_\alpha(p)]. \end{aligned} \quad (28)$$

This expression has a clear physical meaning; $\Delta_4(p)$, the four-dimensional massless propagator, corresponds to the zero-mode graviton localized on the brane, while

$$\Delta_{KK}(p) = -\frac{1}{p} \frac{K_0(pL)}{K_1(pL)} \quad (29)$$

is the propagator for the continuum Kaluza-Klein graviton modes. Comparing the first term of (28) to (16), we obtain the relation between four- and five-dimensional Newton’s constants, $G = 2G_5/L = 1/(8\pi LM^3)$ given above. Note that in the above we have taken the brane to be at the end of a single patch of AdS₅, as was done in [8,9]. This corresponds to the case at hand, since the AdS/CFT relations we have employed above pertain to a single copy of AdS₅.

The continuum graviton modes give rise to corrections to the Newtonian potential. At large distances, corresponding to $pL \ll 1$, a small argument expansion for Bessel functions yields

$$\Delta_{KK}(p) = \frac{L}{2} \left(\ln \frac{p^2 L^2}{4} + 2\gamma \right) + \mathcal{O}(p^2), \quad (30)$$

and, just as in (17), is the source of the $1/r^3$ correction to the Newtonian potential. For a static gravitational source of mass m on the brane, $T_{00}(p) = 2\pi\delta(p_0)m$, evaluating the Fourier transform for $r \gg L$ yields the linearized metric [16]

$$\begin{aligned} h_{00} &= \frac{2Gm}{r} \left(1 + \frac{2L^2}{3r^2} + \dots \right), \\ h_{ij} &= \frac{2Gm}{r} \left(1 + \frac{L^2}{3r^2} + \dots \right) \delta_{ij}, \end{aligned} \quad (31)$$

from which one may read off the Newtonian potential (5).

Moreover, all the metric components in (31) agree with those of (18) and not merely the g_{00} component. In momentum space, this may be traced to the behavior of $h_{\mu\nu}$ in the two pictures, namely (16) and (28). In (28) the factor of $-1/3$ in the non-leading term, as compared with factor $-1/2$ in the leading term, is attributable to the fact that the Kaluza-Klein gravitons are massive. Whereas in (16), it is because the CFT requires loop corrections with $\Pi_2(p) = -\frac{3}{2}\Pi_1(p)$, which is in fact satisfied, as far as the $\ln p^2$ term is concerned, since $a_2 = -\frac{3}{2}a_1$.

We have thus demonstrated that the $1/r^3$ corrections to Newton's law are identical between the Maldacena and Randall-Sundrum pictures. This was examined in the context of a single CFT corresponding to a one-sided brane-world scenario. Had we chosen instead to take the brane-world to be sitting between two patches AdS_5 (one on either side), as was the case considered in [14,10], we would have obtained a factor of two in the relation between Newton's constants, with a corresponding factor in the propagator, (28). While this would ensure the correct four-dimensional behavior of gravity, given in (31), the two-sided brane-world relation $G = G_5/L$ will modify the comparison with the one-loop CFT result, (24). To compensate for this mismatch, one may assume that the two-sided brane-world is dual to two copies of the CFT coupled to gravity, as is implicit in [10]. This leads to the natural picture that a one-sided brane corresponds to a single CFT while a two-sided brane corresponds to two CFTs.

An intriguing feature of this comparison of the gravitational potential in both pictures is a highlighting of the classical/quantum nature of this duality, as seen in the relation

$$\Pi_2(p) + \mathcal{O}(G^2) = \frac{L}{4} \Delta_{KK}(p). \quad (32)$$

The propagator for the continuum graviton modes in the Randall-Sundrum picture thus incorporates all quantum effects of matter on the brane. It may be worthwhile to examine this relation at the two-loop or higher level. Nevertheless, this agreement at one-loop lends strong support to the conjectured duality between the two pictures.

We would like to acknowledge the referees for providing valuable comments. JTL wishes to thank I. Giannakis and H.C. Ren for fruitful discussions on linearized gravity in the brane-world.

REFERENCES

- [1] M. J. Duff, *Problems in the classical and quantum theories of gravitation*, Ph. D. thesis, Imperial College, London (1972).
- [2] M. J. Duff, *Quantum corrections to the Schwarzschild solution*, Phys. Rev. **D9**, 1837 (1974).
- [3] D. M. Capper, M. J. Duff and L. Halpern, *Photon corrections to the graviton propagator*, Phys. Rev. **D10**, 461 (1974).
- [4] D. M. Capper and M. J. Duff, *One-loop neutrino corrections to the graviton propagator*, Nucl. Phys. **B82**, 147 (1974).
- [5] D. M. Capper, *On quantum corrections to the graviton propagator*, Nuovo Cimento **A25**, 29 (1975).
- [6] H. Verlinde, *Holography and compactification*, hep-th/9906182.
- [7] The complementarity and the Newton's law check were suggested independently by E. Witten and J. Maldacena in unpublished work.
- [8] S. S. Gubser, *AdS/CFT and gravity*, hep-th/9912001.
- [9] S. B. Giddings, E. Katz and L. Randall, *Linearized gravity in brane backgrounds*, hep-th/0002091.
- [10] S. W. Hawking, T. Hertog and H. S. Reall, *Brane new world*, hep-th/0003052.
- [11] J. Maldacena, *The large N limit of superconformal field theories and supergravity*, Adv. Theor. Math. Phys. **2**, 231 (1998), hep-th/9711200.
- [12] E. Witten, *Anti-de Sitter space and holography*, Adv. Theor. Math. Phys. **2**, 253 (1998), hep-th/9802150.
- [13] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, *Gauge theory correlators from non-critical string theory*, Phys. Lett. **B428**, 105 (1998), hep-th/9802109.
- [14] L. Randall and R. Sundrum, *An alternative to compactification*, Phys. Rev. Lett. **83**, 4690 (1999), hep-th/9906064.
- [15] L. Susskind and E. Witten, *The Holographic Bound in Anti-de Sitter Space*, hep-th/9805114.
- [16] J. Garriga and T. Tanaka, *Gravity in the Randall-Sundrum Brane World*, Phys. Rev. Lett. **84**, 2778 (2000), hep-th/9911055.
- [17] W. Mück, K. S. Viswanathan and I. V. Volovich, *Geodesics and Newton's law in brane backgrounds*, hep-th/0002132.
- [18] D. M. Capper and M. J. Duff, *Trace anomalies in dimensional regularization*, Nuovo Cimento **A23**, 173 (1974).
- [19] S. Deser, M. J. Duff and C. J. Isham, *Non-local conformal anomalies*, Nucl. Phys. **B11**, 45 (1976).
- [20] M. J. Duff, *Observations on conformal anomalies*, Nucl. Phys. **B125**, 334 (1977).
- [21] M. J. Duff, *Twenty years of the Weyl anomaly*, Class. Quantum. Grav. **11**, 1387 (1994), hep-th/9308075.
- [22] M. Henningson and K. Skenderis, *The holographic Weyl anomaly*, JHEP **9807**, 023 (1998), hep-th/9806087.